

AN IMPLICIT NUMERICAL SCHEME FOR FRACTIONAL ADVECTION DIFFUSION EQUATION

S. M. Chithra, V. Kamalakannan

Department of Mathematics, R.M.K College of Engineering and Technology, Chennai, Tamil Nadu, India

ABSTRACT

In this paper, a finite difference scheme is presented for time fractional advection diffusion equation (TFADE). This equation is derived from classical advection diffusion equation with variable coefficients on replacing classical integer order derivatives by their fractional counterpart. An advection diffusion equation describes physical phenomenon where particle, energy or other physical quantities are transferred inside a physical system due to combined effect of advection and diffusion. To address anomalous diffusion like sub diffusion or super diffusion, classical integer derivatives are replaced by corresponding fractional derivative to obtain TFADE. Using central difference approximations for both space and time fractional derivatives, it is found the present numerical scheme is unconditionally stable.

Keywords: *Finite Difference Method, Advection-Diffusion Equation, Fractional Derivatives.*

I. INTRODUCTION

Fractional Calculus has find its applications in various field of sciences and engineering viz. fluid dynamics, hydrology, acoustic field etc. along with commercial fields such as finance and marketing. Differential equations of arbitrary order (i.e. fractional order) had been used widely to model various systems, material processes, physical and chemical phenomenon. Apart from integer order derivatives, fractional order derivatives address various anomalous phenomenon like sub diffusion and super diffusion. Many attempts are made to find out analytical as well as numerical solution of fractional partial differential equations (FPDE) governing these phenomenon. Among these, sub diffusion and super diffusion processes occurring due to fractional derivative, are studied by many authors (e.g. [2], [3], [10], [12], [14], [15]). Langland and Henry [8] developed numerical algorithm for time fractional

diffusion equation of the form $\frac{\partial^\alpha u(x,t)}{\partial t^\alpha} = \frac{\partial^\alpha u(x,t)}{\partial x^2}$. Considering Riemann Liouville fractional derivative,

Yuste [13] proposed weighted average finite difference method for force free diffusion equation represented by $\frac{\partial u(x,t)}{\partial t} = K_0 D_t^{1-\gamma} \frac{\partial^\alpha u(x,t)}{\partial x^2}$. For $0 < \alpha < 1$, Murio [11] developed unconditionally stable fully implicit

scheme for time fractional diffusion equation of the form $\frac{\partial^\alpha u(x,t)}{\partial t^\alpha} = \frac{\partial^\alpha u(x,t)}{\partial x^2}$ in the space domain $0 < x < 1$

Ding and Li [5] presented a class of weighted finite difference methods for numerical solution of space fractional diffusion equation. Spectral methods based numerical scheme was presented by Lin and Xu [9]. Liu and Yang [10] proposed a numerical method for anomalous sub diffusion equation having a non linear source term. Zhang et.al [14] presented an unconditionally stable finite difference scheme for space time fractional convection diffusion

equation of the form $\frac{\partial^\alpha u(x,t)}{\partial t^\alpha} = a(x) \cdot \frac{\partial^\beta u(x,t)}{\partial t^\beta} - b(x) = \frac{\partial u(x,t)}{\partial x} - c(x)u(x,t) + q(x,t)$ with with initial

and boundary conditions given by $u(0,t)=u(L,t)=0$. Meerschaert et.al. [12] presented finite difference scheme for fractional equation on two time scales. Zhang et.al [15] carried out estimation of error by using fully discrete scheme based on reduction of order. By transforming partial differential equations into an equivalent partial integro differential equation, Huang et.al. [6] constructed two finite difference schemes for time fractional diffusion wave equations. Zhi et.al [4] discussed numerical scheme for one dimensional fractional advection dispersion equation of

the form $\frac{\partial C}{\partial t} + U \frac{\partial C}{\partial x} = K \frac{\partial^2 C}{\partial x^2}$. Yang et.al. [6] proposed a meshless implicit method for 2D time dependent

fractional diffusion wave equation. Using fractional shifted Legendre polynomial method, hybrid numerical scheme

was proposed by Krishnveni et.al [7]. By using an approximation to Reisz fractional derivative, Arshad et.al.[1] presented a finite difference scheme to obtain the numerical solution of time space fractional advection diffusion

equation of the form $C_{D_t^\alpha} u(x,t) = K_{\beta_1} \frac{\partial^{\beta_1}}{\partial |x|^{\beta_1}} u(x,t) + K_{\beta_2} \frac{\partial^{\beta_2}}{\partial |x|^{\beta_2}} u(x,t) + f(x,t)$. The advection

diffusion equation is generally used to model many physical, chemical biological, oceanographic and meteorological

processes. The general form of the advection-diffusion equation is $\frac{\partial u}{\partial t} + v \frac{\partial u}{\partial x} = \frac{\partial^2 u}{\partial x^2}$, where $u(x,t)$ represents

scalar fields like temperature or concentration, $v(x,t)$ represents advective fluid velocity and K is diffusion constant. In the present paper, unconditionally stable fully implicit numerical scheme is presented for time fractional advection diffusion equation (TFADE) of the form

$$\frac{\partial^\alpha u(x,t)}{\partial t^\alpha} + a \frac{\partial u(x,t)}{\partial x} = \frac{\partial}{\partial x} \left(v(x,t) \frac{\partial}{\partial x} u(x,t) \right) \quad (1)$$

On simplification, (1) gets converted into

$$\frac{\partial^\alpha u(x,t)}{\partial t^\alpha} = \frac{\partial v(x,t)}{\partial x} \frac{\partial u(x,t)}{\partial x} + v(x,t) \frac{\partial^2 u(x,t)}{\partial x^2} - a \frac{\partial u(x,t)}{\partial x} \quad (2)$$

To the best of our knowledge no numerical scheme is proposed for this form of advection diffusion equation. In the present scheme, space domain is restricted to $[0,1]$. Along with initial conditions, Dirichlet Boundary conditions are used.

II. NUMERICAL SOLUTION OF TFADE

This section deals with the numerical approximation of TFADE using Caputo fractional time derivative order $0 < \alpha < 1$ given by

$$\frac{\partial^\alpha u(x,t)}{\partial t^\alpha} = \frac{1}{\Gamma(n-\alpha)} \int_0^t \frac{\partial u(x,t)}{\partial t} (t-z)^{-\alpha} dz, \quad 0 \leq t \leq T, \quad 0 < \alpha < 1$$

Let h and k be the mesh size in space and time respectively such that $h=1/M$ and $k=1/N$, where M, N represent number of subintervals in space and time partition respectively. With these implications spatial grid points are given by $x_j = jh$ where $j=0,1,\dots,M$ and grid points in time interval $[0,1]$ are given by $t_r = rk$ where $r = 0,1,\dots,TN$.

Numerical approximation of the solution of TFADE at each grid point is given by $u_j^r = u(x_j, t_r)$ whereas the value of function f at (x_j, t_r) is denoted by $f_j = f(x_j)$. Now we discretize (2) on grid points (x_j, t_r) , we have

$$\frac{\partial^\alpha u(x_j, t_r)}{\partial t^\alpha} = \frac{\partial v(x_j, t_r)}{\partial x} \frac{\partial u(x_j, t_r)}{\partial x} + v(x_j, t_r) \frac{\partial^2 u(x_j, t_r)}{\partial x^2} - a \frac{\partial u(x_j, t_r)}{\partial x} \quad (3)$$

Using numerical approximation to time fractional derivative of first order as suggested by Murio [11] we have

$$\frac{\partial^\alpha u(x_j, t_r)}{\partial t^\alpha} = D_t^\alpha u_j^r + O(k) \quad (4)$$

Where

$$D_t^\alpha u_i^n = \sigma_{\alpha,k} \sum_{r=1}^{r=n} w_r^\alpha (u_i^{n-r+1} - u_i^{n-r}) \quad \text{and} \quad \sigma_{\alpha,k} = \frac{1}{(1-\alpha)\Gamma(1-\alpha)} \frac{1}{k^\alpha} \quad (5)$$

Using central difference approximations to $\frac{\partial u}{\partial x}$, $\frac{\partial v}{\partial x}$, $\frac{\partial^2 u}{\partial x^2}$, we have

$$\frac{\partial v(x_j, t_r)}{\partial x} = \frac{v(x_{j+1}, t_r) - v(x_{j-1}, t_r)}{2h} + O(h^2) \quad (6)$$

$$\frac{\partial u(x_j, t_r)}{\partial x} = \frac{u(x_{j+1}, t_r) - u(x_{j-1}, t_r)}{2h} + O(h^2) \quad (7)$$

$$\frac{\partial^2 u(x_j, t_r)}{\partial x^2} = \frac{u(x_{j-1}, t_r) - 2u(x_j, t_r) + u(x_{j+1}, t_r))}{h^2} + O(h^2) \quad (8)$$

If U_j^r and V_j^r denote the numerical approximations of $u(x_j, t_r)$ and $v(x_j, t_r)$ respectively, then substituting equations (6), (7) and (8), equation (3) reduces to

$$\begin{aligned} \sigma_{\alpha,k} \sum_{r=1}^{r=n} w_r^{(\alpha)} (U_j^{n-r+1} - U_j^{n-r}) &= \left[\frac{V_{j+1}^r - V_{j-1}^r}{2h} \right] \left[\frac{U_{j+1}^r - U_{j-1}^r}{2h} \right] + V_j^r \left[\frac{U_{j-1}^r - 2U_j^r + U_{j+1}^r}{h^2} \right] \\ &\quad - a \left[\frac{U_{j+1}^r - U_{j-1}^r}{2h} \right] \\ &= \left[\frac{V_{j+1}^r - V_{j-1}^r}{4h^2} + V_j^r - \frac{a}{2h} \right] U_{j+1}^r + \left[\frac{V_{j-1}^r - V_{j+1}^r}{4h^2} + V_j^r - \frac{a}{2h} \right] U_{j-1}^r - 2V_j^r U_j^r \end{aligned}$$

With boundary conditions $U_0^n = U_M^n = 0$ and initial conditions $U_j^0 = f_j$, for $j = 1, 2, \dots, M-1$

III. STABILITY ANALYSIS OF NUMERICAL SCHEME

For stability analysis, equation is reconstituted as follows

$$\begin{aligned} \sigma_{\alpha,k} \sum_{r=1}^{r=n} w_r^{(\alpha)} (U_j^{n-r+1} - U_j^{n-r}) &= \left[\frac{V_{j+1}^r - V_{j-1}^r}{4h^2} + V_j^r - \frac{a}{2h} \right] U_{j+1}^r + \left[\frac{V_{j+1}^r - V_{j-1}^r}{4h^2} - V_j^r - \frac{a}{2h} \right] U_{j-1}^r \\ &\quad - 2V_j^r U_j^r \\ \sigma_{\alpha,k} \sum_{r=1}^{r=n} w_r^{(\alpha)} (U_j^{n-r+1} - U_j^{n-r}) &= \left[\frac{V_{j+1}^r - V_{j-1}^r}{4h^2} + V_j^r - \frac{a}{2h} \right] (U_{j+1}^r - U_{j-1}^r) - 2V_j^r (U_j^r - U_{j-1}^r) \end{aligned}$$

If $n=1$, then $r=1$ and we have

$$\sigma_{\alpha,k}(U_j^1 - U_j^0) = \left[\frac{V_{j+1}^1 - V_{j-1}^1}{4h^2} + V_j^1 - \frac{a}{2h} \right] (U_{j+1}^1 - U_{j-1}^1) - 2V_j^1 (U_j^1 - U_{j-1}^1)$$

For $n \geq 2$

$$\sigma_{\alpha,k} \sum_{r=1}^{r=n} w_r^{(\alpha)} (U_j^{n-r+1} - U_j^{n-r}) = V_{j,r} (U_{j+1}^r - U_{j-1}^r) - 2V_j^r (U_j^r - U_{j-1}^r)$$

Where

$$V_{j,r} = \left[\frac{V_{j+1}^r - V_{j-1}^r}{4h^2} + V_j^r - \frac{a}{2h} \right]$$

$$\sigma_{\alpha,k} (U_j^r - U_j^{r-1}) + \sigma_{\alpha,k} \sum_{r=2}^{r=n} w_r^{(\alpha)} (U_j^{n-r+1} - U_j^{n-r}) = V_{j,r} (U_{j+1}^r - U_{j-1}^r) - 2V_j^r (U_j^r - U_{j-1}^r)$$

If we put $U_p^n = \xi_n e^{iwp h}$, where $i = \sqrt{-1}$ and w is a real number, then equation (7) becomes

$$\begin{aligned} \sigma_{\alpha,k} (\xi_n e^{iwp h} - \xi_{n-1} e^{iwp h}) + \sigma_{\alpha,k} \sum_{r=2}^{r=n} w_r^{(\alpha)} (\xi_{n-r+1} e^{iwp h} - \xi_{n-r} e^{iwp h}) &= V_{j,r} (\xi_r e^{i w(p+1)h} - \xi_r e^{i w(p-1)h}) \\ &\quad - 2V_j^r (\xi_r e^{iwp h} - \xi_r e^{i w(p-1)h}) \end{aligned}$$

Equating real part and cancelling $e^{iwp h}$ throughout we have

$$\sigma_{\alpha,k} (\xi_r - \xi_{r-1}) + \sigma_{\alpha,k} \sum_{r=2}^{r=n} w_r^{(\alpha)} (\xi_{n-r+1} - \xi_{n-r}) = -2V_j^r \xi_r (1 - \cos wh)$$

On further simplifications, the above equation can be reduced to

$$\xi_r = \frac{\sigma_{\alpha,k} \left[\xi_{r-1} + \sum_{r=2}^{r=n} w_r^{(\alpha)} (\xi_{n-r} - \xi_{n-r+1}) \right]}{\sigma_{\alpha,k} + 2V_j^r (1 - \cos wh)}$$

or

$$\xi_r = \frac{\xi_{r-1} + \sum_{r=2}^{r=n} w_r^{(\alpha)} (\xi_{n-r} - \xi_{n-r+1})}{1 + \frac{2V_j^r}{\sigma_{\alpha,k}} (1 - \cos wh)}$$

Since $V(x, t) \geq 0$ and $-1 \leq \cos wh \leq 1$, $1 + \frac{2V_j^r}{\sigma_{\alpha,k}} (1 - \cos wh) \geq 1$. Now equation (9) suggests that

$$\xi_1 \leq \xi_0 \text{ and for } r \geq 2$$

$$\xi_r \leq \xi_{r-1} + \sum_{r=2}^{r=n} w_r^{(\alpha)} (\xi_{n-r} - \xi_{n-r+1})$$

Since for each r , $w_r^{(\alpha)} \geq 0$, $w_r^{(\alpha)}(\xi_{n-r} - \xi_{n-r+1}) \leq w_r^{(\alpha)}\xi_{n-r}$, we have

$$\xi_r \leq \xi_{r-1} + \sum_{r=2}^{r=n} w_r^{(\alpha)} \xi_{n-r}$$

If $r=2$, then

$$\begin{aligned} \xi_2 &\leq \xi_1 + w_2^{(\alpha)} \xi_0 \\ &\leq \xi_0 + w_2^{(\alpha)} \xi_0 \\ &= (1 + w_2^{(\alpha)}) \xi_0 \\ &= k_2 \xi_0, \quad \text{where } k_2 = 1 + w_2^{(\alpha)} \end{aligned}$$

If $r=3$, then

$$\begin{aligned} \xi_3 &\leq \xi_2 + w_2^{(\alpha)} \xi_1 + w_3^{(\alpha)} \xi_0 \\ &\leq k_2 \xi_0 + w_2^{(\alpha)} \xi_0 + w_3^{(\alpha)} \xi_0 \\ &= (k_2 + w_2^{(\alpha)} + w_3^{(\alpha)}) \xi_0 \\ &= k_3 \xi_0, \quad \text{where } k_3 = k_2 + w_2^{(\alpha)} + w_3^{(\alpha)} \end{aligned}$$

Continuing like this, we have $\xi_n \leq k_n \xi_0$. Let $k = \max\{1, k_1, k_2, k_3, \dots, k_n\}$. Then $\xi_n \leq k \xi_0$. This shows that the numerical solution is bounded by initial conditions which in turns asserts unconditional stability.

IV. CONCLUSION

The numerical solution of time fractional advection diffusion equation is computed by taking space domain $[0,1]$ and Dirichlet boundary conditions. On the basis of theoretical computations, it is established that the proposed scheme is fully implicit and unconditionally stable.

REFERENCE

1. C.M. Chen, F.Liu, I. Turner, V. Anh, "A Fourier method for the fractional diffusion equation describing sub-diffusion". *Journal of Computational Physics* 2007, 227(2), 886-897.
2. C.M. Chen, F.Liu, I. Turner, V. Anh, "Numerical methods for solving a two dimensional variable-order anomalous subdiffusion equation". *Mathematics of Computation* 2012, 81, 345-366.
3. D. A. Murio, "Implicit finite difference approximation for time fractional diffusion equations". *Computers & Mathematics with Applications* 2008, 56 (4), 1138-1145.
4. F. Liu, C. Yang, K. Burrage, "Numerical method and analytical technique of the modified anomalous subdiffusion equation with a nonlinear source term". *Journal of Computational and Applied Mathematics* 2009, 231(1), 160-176.
5. J. Huang, Y. Tang, L. Vazquez, J. Yang, "Two finite difference schemes for time fractional diffusion-wave equation". *Numerical Algorithms* 2013, 64 (4), 707-720.
6. K. Krishnaveni, K. Kannan, S. R. Balachandar, S.G., Venkatesh, "An efficient fractional polynomial method for space fractional diffusion equations". *Ain Shams Engineering Journal* 2018, 9(4), 767-776.
7. M.M. Meerschaert, Y. Zhang, B. Baeumer, "Particle tracking for fractional diffusion with two time scales". *Computers & Mathematics with Applications* 2010, 59(3), 1078-1086.
8. P. Zhuang, F. Liu, V. Anh, I. Turner, "Stability and convergence of an implicit numerical method for the non-linear fractional reaction-subdiffusion process". *IMA Journal of Applied Mathematics* 2009, 74(5), 645-667.
9. S. Arshad, D. Baleanu, J. Huang, M. M. Qurashi, Y. Tang, Y. Zhao, "Finite difference method for time-space fractional advection diffusion equations with Riesz derivative". *Entropy* 2018, 321, 20(5), 1-20.

10. S.B. Yuste, "Weighted average finite difference methods for fractional diffusion equations. *Journal of Computational Physics* 2006, 216(1), 264-274.
11. T. A. M. Langlands, B.I. Henry, "The accuracy and stability of an implicit solution method for the fractional diffusion equation". *Journal of Computational Physics* 2005, 205, 719-736 (2005).
12. Y. Lin, C. Xu, "Finite difference/spectral approximations for the time-fractional diffusion equation". *Journal of Computational Physics* 2007, 225, 1533-1552.
13. Y. N. Zhang, Z.Z. Sun, H.W. Wu, "Error estimates of Crank-Nicolson-type difference schemes for the subdiffusion equation". *SIAM Journal on Numerical Analysis* 2011, 49(6) 2302-2322.
14. Z. Ding, A. Xiao, M. Li, "Weighted finite difference methods for a class of space fractional partial differential equations with variable coefficients". *Journal of Computation and Applied Mathematics* 2010, 233, 1905-1914.
15. Z.Q. Deng, V.P. Singh, L. Bengtsson, "Numerical solution of fractional advection dispersion equation". *Journal of Hydraulic Engineering* 2004, DOI: 10.1061/(ASCE)0733-9429 (2004) 130:5, 422-431.